

## **A Reanalysis of the 1921 to 1930 Atlantic Hurricane Database**

Christopher W. Landsea\*, Steve Feuer\*, Andrew Hagen\*\*, David A. Glenn\*\*\*, Nicholas T. Anderson\*\*, James Sims\*\*\*\*, Ramon Perez<sup>&</sup>, and Michael Chenoweth<sup>&&</sup>

\*NOAA/NWS/NCEP/National Hurricane Center, Miami, Florida, USA; \*\*University of Miami, Miami, Florida, USA; \*\*\*NOAA/NWS/WFO Morehead City, Morehead City, North Carolina, USA; \*\*\*\*NOAA/NWS/NCEP/Environmental Modeling Center, Camp Springs, Maryland, USA, <sup>&</sup>Institute of Meteorology, Havana, Cuba; <sup>&&</sup>Independent Scholar, Elkridge, Maryland

Submitted to *Journal of Climate*

May 2011

## ABSTRACT

A reanalysis of the Atlantic basin tropical storm and hurricane database ("best track") for the period of 1921 to 1930 has been completed. This reassessment of the main archive for tropical cyclones of the North Atlantic Ocean, Caribbean Sea and Gulf of Mexico was necessary to correct systematic biases and random errors in the data as well as to search for previously unrecognized systems. Methodology for the reanalysis process for revising the track and intensity of tropical cyclone data has been detailed in a previous paper on the reanalysis. The 1921-1930 dataset now includes several new tropical cyclones, excludes one system previously considered a tropical storm, makes generally large alterations in the intensity estimates of most tropical cyclones (both toward stronger and weaker intensities), and typically adjusts existing tracks with minor corrections. Average errors in intensity and track values are estimated for both open ocean conditions as well as for landfalling systems. Finally, highlights are given for changes to the more significant hurricanes to impact the United States, Central America and the Caribbean for this decade.

## **1. Introduction**

This paper details efforts to re-analyze the National Hurricane Center's (NHC's) North Atlantic hurricane database (or HURDAT), also called the “best track” since they are the “best” post-season determination of tropical cyclone (TC) tracks and intensities for the period of 1921 to 1930. Previous work on the reanalysis which have been officially included into the HURDAT data set includes the period of 1851 to 1910 (Landsea et al. 2004a), 1911 to 1920 (Landsea et al. 2008), and 1992's Hurricane Andrew (Landsea et al. 2004b). As the methodology and observational data are nearly identical to that reported for the 1911 to 1920 reanalysis efforts, this short note focuses upon the reanalysis results for the decade of 1921 to 1930. The reader is referred to Landsea et al. (2008) for discussion of data sets utilized and methodology employed.

## **2. New Datasets and Methodology**

The limited observational capabilities of the 1920s were quite similar to that of the previous few decades: measurements from unfortunately placed ships at sea and from coastal weather stations (Landsea et al. 2004a, 2008). The one significant change was with regards to the type of anemometer utilized in United States based observing sites.

The original four-cup anemometer, first developed by Robinson in the 1840s (Kinsman 1969), was still widely used in the United States until the 1920s and in other countries for some time after that. Its primary limitations were in calibrating the instrument and its mechanical failure in hurricane-force wind conditions. Even as late as the 1890s, the highest wind that could be reliably calibrated with this instrument was only about 30 kt (from a whirling machine – similar in structure to a record player), due to a lack of reliable comparisons with a known quantity of faster

motion. By the early 1920s, wind tunnels allowed for calibration against much stronger winds. These showed that the winds from the early cup anemometers had a strong overestimation bias, which was most pronounced at hurricane-force wind speeds (Fergusson and Covert 1924, Kadel 1926). For example, when these instruments indicated winds of minimal hurricane-force of 64 kt, the true wind was only 50 kt. Moreover, most of these early four-cup anemometers were disabled or destroyed by the TC before sampling the highest winds. One of the strongest observed winds in an Atlantic hurricane by this type of anemometer was a 5-min peak wind measurement of 100 kt in Storm #7, 1926 (“The Great Miami Hurricane”) at the U.S. Weather Bureau station in Miami, Florida. (A standard of 5-min was typically utilized in U.S. Weather Bureau reports of “maximum winds” in the era, due to instrumental uncertainties in obtaining shorter time period winds.) With the availability of reliable calibrations beginning in the 1920s, the true velocity of this observation was determined to be only about 77 kt. Current understanding of gustiness in hurricane conditions suggest a boost of 1.06 to convert from a 5-min to a 1-min maximum wind (Powell et al. 1996), giving a best estimate of the maximum 1-min wind of about 82 kt. These older style anemometers were replaced by the more reliably calibrated three-cup anemometers by January 1928 (U.S. Weather Bureau 1927), though these new instruments still suffered from mechanical failure in extreme winds. Anemometers from other countries were gradually upgraded to the new three-cup style generally during the next decade. These corrections to the older style anemometer were thus applied through 1927 for U.S. Weather Bureau stations and for the entire decade elsewhere.

Methodology for re-examining the existing track, intensity, and classification of TCs, for uncovering previously unidentified TCs, and for potentially removing TCs from the database is detailed in Landsea et al. (2008) and is unchanged for what was utilized here for 1921 to 1930. However, new work by Vickery et al. (2009) did allow for a more rigorously defined adjustment of

hurricane winds when converting from marine to open terrain exposure (roughness lengths of 0.03 m). They indicated that a 15% reduction in 1-min winds is appropriate. This is applied in the somewhat rare occasions when a central pressure was available for a TC after landfall, but no direct measurement of the peak winds in the system. In this case, the central pressure was converted to a maximum marine exposure wind speed (Brown et al. 2006) and then the 15% reduction was then applied. Before the availability of the Vickery et al. (2009) paper, a rough reduction of about the same amount was assumed.

One new secondary source for historical hurricanes is Neely (2006) for the country of the Bahamas. This book helped to provide impact information for hurricanes in the Bahamas, as little quantitative observations were typically available from this country during this decade.

### **3. Track, Intensity and Frequency Error Estimates**

Given that the observational datasets for TCs during 1921 to 1930 were nearly the same as for previous decades and that the methodology for reanalysis had not changed, estimates for errors and biases are unchanged from the previous decades (Table 1). The estimated average position errors do depend on whether the TC was out over the open ocean or making landfall, the former being significantly uncertain (~100 nm) and the latter more accurate (~60 nm). (Before the 1920s, some coastal areas were not sufficiently monitored to allow for a more accurate assessment of position compared with the open ocean cases – Landsea et al. [2008].) It is estimated that the intensity measurements for 1921 to 1930 were in error an average of 20 kt over the open ocean, with a bias toward underestimating the true intensity (Table 1). For TCs landfalls during the 1920s, errors in the intensity estimates are smaller - ~12 kt – and likely have a negligible bias as nearly all coastlines around the western North Atlantic, Gulf of Mexico and Caribbean Sea were substantially settled and monitored by then. These estimated errors are the same as the preceding

couple of decades. Landsea et al. (2008a) has additional information on the position and intensity error estimates for the reanalysis database relevant for this decade.

Figure 1 provides an analysis that explicitly quantifies an estimate of what the low bias is for open ocean TCs in the late 19<sup>th</sup> and early 20th centuries. The figure shows in green the hurricanes intensities at the time of impact in the continental U.S. with means of both 89 kt for 1851 to 1930 and for 1980 to 2009. The red bars indicate the difference between the US landfall intensity and the peak lifetime intensity of the hurricane, which is 5.0 kt for 1851 to 1930 and 17.8 kt for 1980 to 2009. For the earlier period, 73% (83 of 113) experienced their peak intensity – according to the reanalyzed HURDAT – right at the time of U.S. landfall, whereas only 38% (17 of 45) of the recent U.S. hurricanes experienced their peak at the time of U.S. landfall. An example of a typical case in the 1980 to 2009 would be Hurricane Katrina in 2005, which peaked over the open Gulf of Mexico as a 150 kt Category 5 on the Saffir-Simpson Hurricane Wind Scale (Landsea et al. 2010), but weakened to a high end Category 3 hurricane with 110 kt maximum 1-min winds to impact the Louisiana coast a day later. Conversely, a typical example for the 1851 to 1930 era would be Storm #3 in 1926 that came ashore in Louisiana with 100 kt maximum 1-min winds, which was also indicated to be the peak intensity for the cyclone. For those more uncommon systems in the earlier period with large (at least 20 kt) differences between the lifetime maximum intensity and the U.S. landfalling intensity, this was almost always due to the hurricane making landfall on another coastline (such as Cuba in Storm #10, 1924) at a higher intensity than that at U.S. landfall.

The quite sizable discrepancy in the peak intensity minus U.S. landfall intensity differences – only 5.0 kt for the earlier era and 17.8 kt for the modern hurricanes – requires an explanation. One possibility is that the populations being sampled from both eras are too dissimilar for a

homogeneous comparison. However, evaluating the characteristics of each suggests that the TCs are being drawn from quite similar samples: mean intensities at U.S. impact: 89 kt - 1980-2009, 89 kt - 1851-1930; % Atlantic coast/% Gulf coast landfalls: 31%/69% - 1980-2009, 37%/63% - 1851-1930; % striking in August-October: 78% - 1980-2009, 80% - 1851-1930. Given that the sample differences do not explain the discrepancy, a likely reason is that the modern era calculations show that most TCs have a real, substantial weakening before U.S. landfall (especially in the Gulf of Mexico for major hurricanes – Rappaport et al. 2010), whereas the inability to adequately monitor the intensities of TC over the open ocean in the 1851 to 1930 era results in not capturing the peak lifetime intensity of TCs. This discrepancy – about 13 kt – is quite similar to the more subjectively determined estimates of 10-15 kt low bias for open Atlantic TC intensity provided in Table 1 and estimated previously in Landsea et al. (2004, 2008a).

Recent research (Chang and Guo 2007, Vecchi and Knutson 2008, Landsea et al. 2010) has allowed for more reliable estimates of the number of “missing” TCs before the advent of satellite imagery. The Vecchi and Knutson (2008) results suggest that there were about one missed tropical cyclone per year during the 1920s. Landsea et al. (2010) also indicated that there has been an extreme increase in the number of short-lived (less than or equal to a two day duration of tropical storm or greater intensity) TCs in the last couple of decades that is likely due to better technology and monitoring of these short-lived and typically very weak systems. Compared to rates of short-lived TCs in recent years, this suggests (Landsea et al. 2010) that about four short-lived TCs were missed per year in the 1920s and about one medium to long-lived TCs (greater than a two day duration) missed every other year. These conclusions will be put into the context of the results of the reanalysis, which did lead to a substantial change in the frequency of all TCs and short-lived TCs.

#### **4. Results:**

##### *a. Overall activity:*

A summary of the yearly changes to HURDAT is provided in Figure 2 and Table 2. Figure 2 shows the revised and comparison track maps for the individual seasons from 1921 to 1930. It is apparent that most of the track changes introduced for these years are fairly minor (less than a 120 nmi alteration in position at anytime during the TC's lifetime) as readily seen in the comparison maps, though there are some more dramatic alterations on occasion (e.g., Storm #5 in 1923, Storm #1 in 1927, and Storm #1 in 1928). Despite making relatively minor changes overall, nearly every existing TC was adjusted for at least some portion of its track.

In addition to track alterations of existing systems, new TCs were discovered and added into HURDAT and one existing system in HURDAT was reanalyzed to not be a tropical storm and thus removed from the database. In total, 14 new TCs had sufficient observational evidence to document their existence and were added into HURDAT: 3 in 1923 and 1924; 2 in 1925 and 1929; 1 in 1921, 1922, 1927, and 1930; and no new systems in 1926 and 1928. Of these 14, two of the new TCs were landfalling systems: Storm #7, 1921 in Cuba and Storm #1, 1923 in Louisiana, Mississippi, and Alabama. Additionally, one system during the 1920s in HURDAT was removed because of a lack of gale force winds (original Storm #4 in 1923). In other years in the reanalysis work (e.g., 1891), two separate TCs were found to be actually one continuous system and thus so changed to reflect this, but no such systems were uncovered during the 1920s.

Table 2 lists the original and revised tallies of tropical storms and hurricanes, hurricanes, major hurricanes (Category 3, 4 and 5 on the Saffir-Simpson Hurricane Wind Scale [SSHWS]),



and Accumulated Cyclone Energy (ACE – an index for overall TC activity that takes into account the total frequency, intensity and duration of TCs, Bell et al. 2000). ACE is calculated by summing the squares of the estimated 6-hourly maximum wind speed in knots to be found in HURDAT for all periods while the system is either a tropical storm or hurricane.

The average number of tropical storms and hurricanes increased from 5.6 per year in the original HURDAT to 6.9 after the reanalysis (Table 2), a 23% increase. This net increase includes new systems that we added into the database as well as the one that was originally in HURDAT but was discarded. Both values are substantially below the long-term average of 11.2 per year recorded in the satellite era of 1966-2009 (updated from Blake et al. 2007). However, as described earlier, a direct comparison of the total frequency of TCs during the 1920s to the modern climatology is complicated by the occurrence of “missed” TCs in the earlier years because of a lack of satellite imagery and vastly improved monitoring capability available now. In the original HURDAT, of the 56 TCs, only three were short-lived. With the reanalysis, of the 69 TCs for the 1921-30 period, 11 are now indicated to be short-lived TCs. Five of the newly described short-lived TCs were due to a decrease in the original duration recorded and four were brand new TCs not previously recorded. To better homogeneously compare the 1920s to the more recent era, one must estimate the number of “missed” TCs of medium to long durations in the 1921-30 period and remove the likely spurious influence of the short-lived TC trends. Using the results of Landsea et al. (2010), an average of about one medium to long-lived TC every two years were missed in the 1920s and the modern (1966 to 2009) climatology of medium to long-lived TCs is 7.7 per year. Thus the best adjusted total of medium to long-lived TCs from 1921 to 1930 is about 6.3 per year, which suggests that this decade was quiet relative to the modern era for TC frequency.

In contrast, the hurricane, major hurricane and ACE averages (Table 2) show smaller changes in recorded values. Hurricane frequency had a small increase from 3.6 to 4.0 per year (6.1 per year in the modern era), major hurricanes remained unchanged at 1.7 per year (2.3 per year recently), and ACE increased slightly from 71.8 to 76.6 per year (94.5 per year recently). With regards to ACE, three years recorded a substantial increase in activity (ACE higher by at least 10.0 – 1921, 1923, and 1930), two years saw a decrease in activity (ACE lower – 1922 and 1923), and the remaining five years had minor increases in overall intensity, duration and frequency. To have ACE with little change or even a modest decrease in some years is likely due to a systematic tendency for the original HURDAT to somewhat overestimate the intensity of hurricanes from 1921 to 1930, especially over the open ocean (e.g., Storm #4, 1927). In general, large changes to intensity (at least a 20 kt alteration at some point in the TC’s lifetime) were recorded – both upward and downward – for the majority of individual TCs, typically with more significant changes than those introduced for track. Currently, there exists no method for quantifying the amount of “missed” hurricanes, major hurricanes, and ACE for the era of the 1920s. Because of this any direct comparison of these quantities to the modern era would not be appropriate.

*b. Continental United States Hurricanes:*

Table 3 summarizes the continental U.S. hurricanes for the period of 1921 to 1930 and the states impacted by these systems. U. S. hurricanes are defined as those hurricanes that are analyzed to cause maximum (1 min) surface (10 m) winds of at least 64 kt for an open exposure on the coast or inland in the continental United States. Hurricanes that make a direct landfall with the circulation center (eye) of the system crossing the coast as well as those that make a close bypass are considered. In addition to the parameters also common to HURDAT (e.g. latitude, longitude,

maximum winds and central pressure), the U.S. hurricane compilation also includes the outer closed isobar, the mean size of the outer closed isobar, and - when available - the radius of maximum wind (RMW). These parameters provide information regarding the size of the hurricanes, which can vary considerably from system to system. For these TCs, winds listed in HURDAT in the last six hourly period before landfall are now consistent with the assigned Saffir-Simpson Hurricane Scale category, which was not the case in the original HURDAT database before the reanalysis efforts. For most U.S. hurricanes of this era, a central pressure observation or estimate was obtained from original sources, which was then used to determine maximum wind speeds through the application of one of the Brown et al. (2006) pressure-wind relationships. In the one case where there was no central pressure value directly available, the estimated winds at landfall were then used via the pressure-wind relationship to back out a reasonable central pressure. In either case, the objective was to provide both an estimate of the maximum wind and a central pressure at landfall for all U.S. hurricanes.

There were 16 U.S. hurricanes (five that were major hurricanes) during the 1921 to 1930 period after the reanalysis. No U.S. hurricanes were recorded in 1922, 1927, or 1930. The total of 16 U.S. hurricanes represents two more hurricanes than the original HURDAT database contained, with Storm #3 in 1924 (causing Category 1 hurricane conditions in North Carolina and Massachusetts) and Storm #10 in 1926 (causing Category 1 hurricane conditions in south Florida) as close bypassing systems. (Originally, these two systems were considered to have caused tropical storm impacts for the U.S. in HURDAT.) For existing U.S. hurricanes, thirteen were unchanged in Category, while one was downgraded by a Category: Storm #1, 1921 from a Category 2 in Texas originally to a Category 1. No original U.S. hurricanes were removed as such

from HURDAT during the 1921-1930 time period. No major hurricanes were either added or removed from the U.S. hurricane list.

Notable hurricanes that affected the continental United States for 1921 through 1930 (Blake et al. 2007) include Storm #10, 1921 Category 3 in southwest Florida; Storm #3 in 1926 in Louisiana; Storm #7, 1926 – the “Great Miami Hurricane” in southeast and northeast Florida and Alabama; Storm #4, 1928 – the “Lake Okeechobee Hurricane” in southeast and northwest Florida; and Storm #2, 1929 in south Florida.

During the period of 1921 to 1930, the first very destructive hurricane to strike the continental United States was Storm #10, 1921, which hit the southwest coast of Florida just north of Tampa. This hurricane was the last direct strike by a major hurricane in the Tampa-St. Petersburg metropolitan area (Blake et al. 2007). This TC was originally listed as a Category 3 with a 952 mb central pressure at landfall and a maximum wind at the last synoptic time before landfall of 90 kt<sup>1</sup>. This central pressure was unchanged along with a moderately-sized RMW of 15 nm giving a maximum 1-min wind estimated at landfall to be about 105 kt, retaining the system as a Category 3 at landfall.

The next major hurricane to affect the United States was not until 1926. Storm #3 was originally estimated to have a central pressure of 955 mb at landfall in Louisiana and maximum winds of 90 kt at the last synoptic time before landfall. After the reanalysis, the central pressure was unchanged and along with a moderately-sized RMW of 20 nm, the winds at landfall were

---

<sup>1</sup> The discrepancy between the Category 3 original assessment for U.S. landfall of this hurricane with the 90 kt winds existing originally in HURDAT is a quite common problem in the existing dataset. Much of the discrepancy is due to reliance primarily upon the central pressure by Taylor and Hebert (1975) to provide the original Saffir-Simpson Hurricane Wind Scale Category at landfall in the United States, while the practice today at the National Hurricane Center and within the reanalysis is to determine the maximum winds at landfall and then let these determine the appropriate Category.

estimated to be 100 kt, retaining the Category 3 hurricane status for this system. Also in 1926, Florida and Alabama were devastated by the “Great Miami Hurricane” with about 372 people killed (Pfof 2003). This system was analyzed to have a central pressure of 935 mb at landfall in southeast Florida with winds of 115 kt at the last synoptic time before landfall. At its second landfall in northwest Florida/Alabama, it originally had a central pressure recorded of 955 mb and maximum winds of 105 kt at the last synoptic time before landfall. It was originally assessed to be Category 4 at landfall in southeast Florida and Category 3 at a second landfall in northwest Florida/Alabama. After reanalysis, it was determined that the central pressure at landfall in Miami was slightly deeper at 930 mb and – along with a moderately-sized RMW of 20 nm – had maximum winds of about 125 kt. The second landfall of the hurricane in northwest Florida/Alabama was reassessed to be slightly different than originally estimated: a central pressure of 955 mb and maximum winds of 100 kt. This retains the Category 4 and Category 3 assessments for southeast Florida and northwest Florida/Alabama, respectively. This large (outer closed isobar of 325 nm radius) and powerful hurricane would cause on the order of \$165 billion in total losses today, given society’s population and infrastructure in place now (Blake et al. 2007). Such destruction would place as the most expensive hurricane – after normalization – in U.S. hurricane history since 1900.

Just two years later, southeast Florida suffered another devastating hurricane strike with the impact of Storm #4 – “The Lake Okeechobee Hurricane”. This system killed on the order of 2500 people (Pfof 2003) and was originally assessed to be a 929 mb central pressure Category 4 hurricane at landfall in Palm Beach with maximum winds of 130 kt at the last synoptic time before landfall. After reanalysis, the 929 mb was retained and partly because of a large RMW – 30 nm - for this central pressure and landfall latitude, the maximum winds were assessed to be 125 kt at

landfall. This retains the hurricane as a Category 4 hurricane at landfall in southeast Florida. If this system were to strike the United States today, it would cause on the order of \$35 billion, making it the 8<sup>th</sup> most expensive hurricane in U.S. history back to 1900 normalized to today's societal vulnerability (Blake et al. 2007).

The last major hurricane to make landfall in the United States was Storm #2 in 1929. This system originally made landfall in the Florida Keys as a Category 3 hurricane with 948 mb central pressure and maximum winds of 90 kt at the last synoptic time before landfall. After reanalysis, the 948 mb central pressure was retained and – along with a large RMW of about 30 nm and a slow, 4 kt translational speed – winds of 100 kt were analyzed at landfall. These maximum winds allow for the system to be retained as a Category 3 hurricane at landfall.

Overall, there were no additions to the U.S. major hurricanes for the years of 1921 to 1930, nor were there any alterations to the peak Category impact at landfall. However, maximum winds in HURDAT were adjusted for all five of these U.S. major hurricanes with one decreased by 5 kt (as was the second landfall of Storm #9, 1926), three increased by 10 kt, and one increased by 15 kt.

*c. Major Hurricanes outside of the Continental U.S.:*

Outside of the continental United States, major hurricanes impacted several locations from 1921 to 1930. Nine separate major hurricanes made landfall either in the Lesser Antilles, Greater Antilles or Bermuda. Of note was that all of Central America including all of the east coast of Mexico was spared from any direct strikes by major hurricanes during this time period. Of the

nine, one was newly designated to be a major hurricane after the reanalysis: Storm #3 in 1921 that struck the Dominican Republic, the winds from which were increased from 80 to 110 kt at landfall.

The largest impact of any major hurricane was the devastating Category 4 hurricane that struck also Dominican Republic in September 1930 (Storm #2, 1930) and killed in the range of 2,000 to 8,000 people (Rappaport and Partagás 1995). For this hurricane – the worst impact and most intense in the Dominican Republic’s history – had an observed 933 mb central pressure and estimated 135 kt maximum winds at landfall.

Cuba was struck by two major hurricanes during this decade: a Category 5 hurricane – the “Huracán sin Precedentes” – in October 1924 (Storm #10, 1924) and a Category 4 hurricane in October 1926 (Storm #10, 1926). The former cyclone caused extraordinary destruction in the western portion of the country (Perez et al. 2000) with an analyzed central pressure of 910 mb and estimated 140 kt maximum winds at landfall. The latter hurricane with an analyzed central pressure of 934 mb and estimated 130 kt maximum winds at landfall killed about 600 people in Cuba (Perez et al. 2000). The “Huracán sin Precedentes” is only one of two Category 5 hurricanes documented to have struck Cuba in its long hurricane history (Perez et al. 2000), the first of which was “La Tormenta de San Francisco de Borja” of October 1846 (preceding the current HURDAT database).

The Bahamas were also struck repeatedly by major hurricanes during the decade with one Category 3 strike (Storm #10, 1926) and four Category 4 strikes (Storm #1, 1926; Storm #7, 1926; Storm #4, 1928; and Storm #2, 1929). Of these Storm #1, 1926 and Storm #2, 1929 had the largest impacts to Bahamian residents with about 300 people killed in the former and extreme destruction caused to Nassau and other islands in the latter (Neely 2006).

Puerto Rico was struck by only one major hurricane during the period of 1921 to 1930: Storm #4, 1928, known as “San Felipe” locally. This system killed over 300 people (Rappaport and Partagás 1995) and is the most intense to ever strike the island with observed central pressure of 931 mb and observed maximum winds of 140 kt. Originally, “San Felipe” was the first Category 5 hurricane to appear in the HURDAT database anywhere within the Atlantic basin (not just at landfall). However, with the upgrade of the “Huracán sin Precedentes” (Storm #10, 1924) to a peak 140 kt intensity, this system has supplanted “San Felipe” as the first Category 5 to appear anywhere within HURDAT.

To summarize significant changes to the landfall intensity of these nine major landfalling (non-continental U.S.) hurricanes, three had large increases in landfall intensity (Storm #3, 1921 from 80 up to 110 kt; Storm #10, 1924 from 105 up to 140 kt; and Storm #10, 1926 from 105 to 130 kt in Cuba), one had a large decrease in landfall intensity (Storm #2, 1922 from 130 down to 100 kt in Bermuda), and the remainder had small or no alterations in landfall intensity.

## **5. Summary and Future Work:**

Continued progress on the hurricane reanalysis has been made through now the 3rd decade of the 20th Century: 1921 to 1930. While the results provided here are just brief summaries of the thousands of changes introduced into the Atlantic hurricane database, all raw observations, the original and revised HURDAT, annual track maps, metadata regarding changes for individual TCs, and comments from/replies to the National Hurricane Center’s Best Track Change Committee can be found at : [http://www.aoml.noaa.gov/hrd/data\\_sub/re\\_anal.html](http://www.aoml.noaa.gov/hrd/data_sub/re_anal.html).

Highlights of accomplishments attained for this stage of the Atlantic hurricane database reanalysis project for 1921 to 1930:



- A) Track alterations were implemented for nearly all TCs in the existing HURDAT, though the majority were for minor changes;
- B) Intensity changes were incorporated into nearly all TCs with a much larger proportion with major alterations in their intensity compared with the track, either toward stronger or weaker winds;
- C) 14 new TCs were discovered and added into HURDAT, while 1 system was removed from the database because it was not of tropical storm intensity;
- D) While the frequency of TCs during the era was increased from 5.6 to 6.9 annually because of these net changes, a smaller increase was noted for hurricane frequency and no net changes were noted in for major hurricane numbers. The overall activity - as denoted by Accumulated Cyclone Energy - was increased by about 7%;
- E) 16 continental U.S. hurricanes were identified, two more than originally listed in HURDAT because of an addition of two new U.S. hurricanes (originally were considered tropical storm impact for the United States) and the removal of none during the time period. Of the 14 original U.S. hurricanes, thirteen had no changes introduced for the peak Category, while one was downgraded a Category. No changes were made to the number of major continental U.S. hurricanes (five) for the decade and none of these were adjusted for their top SSHWS Category impact;
- F) Nine major hurricanes struck other places in the Atlantic basin, one of which was newly classified as a major hurricane. Of the nine, three had large (at least

20 kt) increases of intensity at landfall, while one had a large reduction in landfall winds;

- G) Despite the reanalysis changes, there exists significant uncertainty in TC tracks, significant undercounts in TC frequency, and significant underestimation of TC intensity and duration, especially for those systems over the open ocean.

This last point requires some elaboration. In the recent past, there have been some climate change studies utilizing HURDAT that have assumed essentially no missing TCs or no missing portions of TCs life cycle or no underestimating of the intensity of existing TCs beginning in the mid-20th Century or even back to the beginning of the database in 1851 (e.g., Emanuel 2005, Mann and Emanuel 2006, Holland and Webster 2007). Innovative new work (Vecchi and Knutson 2008, Landsea et al. 2010) has come up with credible, quantitative estimates of how many TCs have been "missed" in the historical database. Such work needs to be extended, if possible, to other TC metrics, such as hurricane and major hurricane frequency, hurricane days, and ACE. A rough measure of the underestimate in the intensity values in HURDAT up to 1930 has been provided in Table 1 with some quantitative analysis in Figure 1 in support of these estimates. It is incumbent upon HURDAT users to not ignore the likely effects of incomplete sampling on creating spurious trends in various TC metrics.

Considerably more work needs to be accomplished for the Atlantic hurricane database. One essential project is to extend HURDAT back before 1851 to earlier in the 19th, 18th and even 17th Centuries. Such efforts are underway and have begun to yield important results (e.g., Chenoweth 2007, Mock 2008, Wheeler et al. 2009). This may lead to a complete dataset of U.S. landfalling hurricanes for the Atlantic coast from Georgia to New England as well as for portions of the Caribbean back to at least 1800, given the relatively high density of population extending that far

into the past. While the reanalysis efforts thus far have extended HURDAT back to 1851 and revised it through 1930, these did not make extensive use of COADS until the decade of the 1910s (Landsea et al. 2004a). Further improvements in HURDAT could be achieved by utilizing this comprehensive ship database for the years of 1851 to 1910. Finally, an ongoing project is to complete the current reanalysis efforts through the remainder of the 20<sup>th</sup> Century. Preliminary results from this research (e.g., Hagen 2010) indicates the potential for much increased accuracy and completeness in HURDAT is possible.

*Acknowledgments.*

This work has been sponsored by a grant from the NOAA Climate Program Office on "Re-analysis of the Atlantic Basin Tropical Cyclone Database in the Modern Era". The authors wish to thank the current NHC Best Track Change Committee (Chair - Jack Beven, Lixion Avila, Eric Blake, Todd, Kimberlain, Richard Pasch, Gladys Rubio, and Eric Uhlhorn) and past members (Hugh Cobb, Jim Gross, Brian Jarvinen, Colin McAdie, and Ed Rappaport) for their encouragement and detailed suggestions that have helped to quality control the thousands of alterations and additions to HURDAT. Special thanks for their individual contributions toward this project for this era are also given to Sim Aberson, James Belanger, Auguste Boissonnade and Risk Management Solutions, Emery Boose, Gil Clark, Daniel Gladstein, Mark Jelinek, Cary Mock, Charlie Neumann, Ricardo Preto, David Roth, Jose Sánchez-Sesma, Al Sandrik, Donna Strahan, and Mark Zimmer. Joan David kindly provided the track map figures utilized in this paper.

## References

Bell, G. D., and Coauthors, 2000: Climate assessment for 1999. *Bull. Amer. Meteor. Soc.*, **81**, S1-S50.

Blake, E. S., E. N. Rappaport, and C. W. Landsea, 2007: The deadliest, costliest, and most intense United States tropical cyclones from 1851 to 2006 (and other frequently requested hurricane facts). *NOAA Technical Memorandum*, NWS-TPC-5, 43 pp. < <http://www.nhc.noaa.gov/pdf/NWS-TPC-5.pdf> >

Brown, D. P., J. L. Franklin, and C. W. Landsea, 2006: A fresh look at tropical cyclone pressure-wind relationships using recent reconnaissance-based “best track” data (1998–2005). Preprints, *27th Conf. on Hurricanes and Tropical Meteorology*, Monterey, CA, Amer. Meteor. Soc., 3B.5. [Available online at <http://ams.confex.com/ams/pdfpapers/107190.pdf>.]

Chang, E. K. M., and Y. Guo, 2007: Is the number of North Atlantic tropical cyclone significantly underestimated prior to the availability of satellite observations? *Geophys. Res. Letters*, **34**, L14801, doi:10.1029/2007GL030169.

Chenoweth, M., 2007: A reassessment of historical Atlantic basin tropical cyclone activity, 1700–1855. *Climatic Change*, **76**, 169–240, doi:10.1007/s10584-005-9005-2.

Emanuel, K. A., 2005: Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, **436**, 686–688.

Fergusson, S. P., and R. N. Covert, 1924: New standards of anemometry. *Mon. Wea. Rev.*, **52**, 216–218.

Hagen, Andrew B., 2010: A reanalysis of the 1944-1953 Atlantic hurricane seasons – The first decade of aircraft reconnaissance. M.S. Thesis, University of Miami, Miami, FL, 851 pp.

<http://etd.library.miami.edu/theses/available/etd-12132010-141954/> .

Hebert, P. J., and G. Taylor, 1975: Hurricane experience levels of coastal county populations from Texas to Maine. National Weather Service Community Preparedness Staff and Southern Region Special Rep., 153 pp.

Holland, G. J. and P. J. Webster, 2007: Heightened tropical cyclone activity in the North Atlantic: natural variability or climate trend? *Philos. Trans. Roy. Soc. A*, **365**, 2695–2716.

Jarvinen, B. R., C. J. Neumann, and M. A. S. Davis, 1984: A tropical cyclone data tape for the North Atlantic Basin, 1886-1983: Contents, limitations, and uses. *NOAA Technical Memorandum NWS NHC 22*, Coral Gables, Florida, 21 pp.

Kadel, B. C., 1926: An interpretation of the wind velocity record at Miami Beach, Fla., September 17-18, 1926. *Mon. Wea. Rev.*, **54**, 414-416.

Kinsman, B., 1969: Who put the wind speeds in Admiral Beaufort's Force Scale? *Oceans*, **2**, 18–25.

Landsea, C. W., C. Anderson, N. Charles, G. Clark, J. Dunion, J. Partagas, P. Hungerford, C.

Neumann and M. Zimmer, 2004a: The Atlantic hurricane database re-analysis project:

Documentation for the 1851-1910 alterations and additions to the HURDAT database. *Hurricanes and Typhoons: Past, Present and Future*, R. J. Murnane and K.-B. Liu, Eds., Columbia University Press, 177-221.

Landsea, C.W., J. L. Franklin, C. J. McAdie, J. L. Beven II, J. M. Gross, R. J. Pasch, E. N.

Rappaport, J. P. Dunion, and P. P. Dodge, 2004b: A reanalysis of Hurricane Andrew's intensity.

*Bull. Amer. Meteor. Soc.*, **85**, 1699-1712.

Landsea, C. W. , D. A. Glenn, W. Bredemeyer, M. Chenoweth, R. Ellis J. Gamache, L. Hufstetler,

C. Mock, R. Perez, R. Prieto, J. Sanchez-Sesma, D. Thomas, and L. Woolcock, 2008: A reanalysis

of the 1911-20 Atlantic hurricane database. *J. Climate*, **21**, 2138-2168.

Landsea, C.W., G.A. Vecchi, L. Bengtsson, and T. R. Knutson, 2010: Impact of duration

thresholds on Atlantic tropical cyclone counts. *J. Climate*, **23**, 2508-2519.

Mann, M. and K. Emanuel, 2006: Atlantic hurricane trends linked to climate change. *EOS*, **87**,

233–241.

Mock, C.J.. 2008: Tropical cyclone variations in Louisiana, U.S.A. since the late Eighteenth

Century. *Geochemistry, Geophysics, and Geosystems*. **9**, Q05V02, doi:10.1029/2007GC001846.

Neely, W., 2006: *The Major Hurricanes to Affect the Bahamas: Personal Recollections of Some of the Greatest Storms to Affect the Bahamas*, AuthorHouse, 252 pp.

Perez Suarez, R., R. Vega, and M. Limia, 2000: *Cronologia de los Ciclones Tropicales de Cuba*.

Instituto de Meteorologia, 100 pp.

Pfost, R. L., 2003: Reassessing the impact of two historical Florida hurricanes. *Bull. Amer.*

*Meteor. Soc.*, **84**, 1367-1372.

Powell, M. D., S. H. Houston, and T. A. Reinhold, 1996: Hurricane Andrew's landfall in south Florida. Part I: Standardizing measurements for documentation of surface wind fields. *Wea. Forecasting*, **11**, 304–328.

Rappaport, E. N., J. L. Franklin, A. B. Schumacher, M. DeMaria, L. K. Shay, and E. J. Gibney, 2010: Tropical cyclone intensity change before U.S. Gulf coast landfall. *Wea. Forecasting*, (in press).

Rappaport, E. N., and J. Fernández-Partagás, 1995: The deadliest Atlantic tropical cyclones, 1492–1994. NOAA Tech. Memo. NWS NHC-47, 41 pp.


U.S. Weather Bureau, 1927: New three-cup anemometer. *Mon. Wea. Rev.*, **55**, 533.

Vecchi, G. A., and T. R. Knutson, 2008: On estimates of historical North Atlantic tropical cyclone activity. *J. Climate*, **21**, 3580–3600.

Vickery, P. J., D. Wadhera, M. D. Powell, and Y. Chen, 2009: A hurricane boundary layer and wind field model for use in engineering applications. *J. App. Meteor. Climatol.*, **48**, 381–405.

Wheeler, D., R. García-Herrera, J. M. Vaquero, M. Chenoweth, and C. J. Mock, 2009: Reconstructing the trajectory of the August 1680 hurricane from contemporary records. *Bull. Amer. Meteorol. Soc.*, **90**, 971–978.

# U.S. Landfalling Hurricane Intensity and Difference from Peak Intensity 1851-1930 - Mean Difference 5.0 kt





## U.S. Landfalling Hurricane Intensity and Difference from Peak Intensity 1980-2008 - Mean Difference 17.8 kt














Figure 1: Comparison of intensity of hurricanes striking the continental United States (in green) versus the difference of that wind speed versus the peak lifetime intensity for that hurricane (in red). Top panel is for 1851 to 1930 and includes only those hurricanes that struck a settled part of the U.S. coast line (Landsea et al. 2004a, 2008), while the bottom panel is for 1980 to 2009 (updated from Blake et al. 2007).

































Figure 2: The revised (top) and comparison (with original track in blue underlying revised track) (bottom) Atlantic basin TC track map for 1921, 1922, 1923, 1924, 1925, 1926, 1927, 1928, 1929, and 1930.

Table 1: Estimated average position and intensity errors in the revised best track for the years 1851-1930. Negative bias errors indicate an underestimation of the true intensity. (By the 1920s, nearly all coastal areas in the Atlantic basin were relatively settled and monitored – Landsea et al. [2008].)

Situation	Dates	Position Error	Intensity Error	Intensity Error
			(absolute)	(bias)
Open ocean	1851-1885	120 nmi	25 kt	-15 kt
	1886-1930	100	20	-10
Landfall at sparsely populated area	1851-1885	120	25	-15
	1886-1920	100	20	-10
Landfall at settled Area	1851-1885	60	15	0
	1886-1930	60	12	0

Table 2: Original/revised tropical storm and hurricane, hurricane, major hurricane and Accumulated Cyclone Energy (ACE) counts. ACE is expressed in units of  $10^4$  kt<sup>2</sup>.

Year	Tropical Storms & Hurricanes	Hurricanes	Major Hurricanes	ACE
1921	6/7	4/5	2/2	75/87
1922	4/5	2/3	1/1	58/55
1923	7/9	3/4	1/1	54/49
1924	8/11	5/5	2/2	89/100
1925	2/4	1/2	0/0	7/8
1926	11/11	8/8	6/6	222/230
1927	7/8	4/4	2/1	56/56
1928	6/6	4/4	1/1	75/83
1929	3/5	3/3	1/1	43/48
1930	2/3	2/2	1/2	39/50
Average 1921-1930	5.6/6.9	3.6/4.0	1.7/1.7	71.8/76.6
Average 1966-2009	11.2	6.1	2.3	94.5

Table 3: Continental U.S. Hurricanes for 1921 to 1930. Date/Time - day and time when the circulation center crosses the U.S. coastline (including barrier islands). Time is estimated to the nearest hour. Lat/Lon - Location is estimated to the nearest 0.1 degrees latitude and longitude. Max Winds - Estimated maximum (1 min) surface (10 m) winds to occur along the U. S. coast. SSHWS - The Saffir-Simpson Hurricane Wind Scale at landfall based upon estimated maximum 1-min surface winds. RMW - The radius of maximum winds, if available, to the nearest 5 nm. Central Pressure - The minimum central pressure of the hurricane at landfall. Central pressure values in parentheses indicate that the value is a simple estimation (based upon a wind-pressure relationship), not directly measured or calculated. OCI – The Outer Closed Isobar, or the sea level pressure at the outer limits of the hurricane circulation as determined by analysis of the outer closed isobar (in increments of 1 mb). Size - This is the average radius of the OCI. States Affected - The impact of the hurricane upon individual U.S. states by Saffir-Simpson Hurricane Wind Scale (again through the estimate of the maximum 1-min surface winds at each state). (ATX-South Texas, BTX-Central Texas, CTX-North Texas, LA-Louisiana, MS-Mississippi, AL-Alabama, AFL-Northwest Florida, BFL-Southwest Florida, CFL-Southeast Florida, DFL-Northeast Florida, GA-Georgia, SC-South Carolina, NC-North Carolina, VA-Virginia, MD-Maryland, DE-Delaware, NJ-New Jersey, NY-New York, PA-Pennsylvania, CT-Connecticut, RI-Rhode Island, MA-Massachusetts, NH-New Hampshire, ME-Maine. In Texas, south is roughly from the Mexico border to Corpus Christi; central is from north of Corpus Christi to Matagorda Bay and north is from Matagorda Bay to the Louisiana border. In Florida, the north-south dividing line is from Cape Canaveral [28.45N] to Tarpon Springs [28.17N]. The dividing line between west-east Florida goes from 82.69W at the north Florida border with Georgia, to Lake Okeechobee and due south along longitude 80.85W.) “\*” - Indicates that the hurricane center did not make a U.S.

landfall, but did produce hurricane force winds over land. Position indicated is point of closest approach. Winds stronger than indicated to impact the United States may have existed elsewhere in the hurricane. Central pressure in this case is the hurricane's value at the point of closest approach.

#/Date	Time	Lat	Lon	Max Winds	SSHWS	RMW	Central Pressure	OCI	Size	States Affected	Original Assessment
1-6/22/1921	1800Z	28.6N	95.9W	80	1	15	980	1011	225	BTX1,CTX1	BTX2
6-10/25/1921	2000Z	28.1N	82.8W	105	3	20	952	1009	375	BFL3,AFL2,DFL1,CFL1	BFL3,DFL2
None - 1922											
6-10/16/1923	0600Z	29.2N	91.0W	70	1	45	983	1001	250	LA1,MS1	LA1
3-8/26/1924*	0400Z	35.5N	74.8W	65	1	35	963	1009	275	NC1	None
3-8/26/1924*	1900Z	41.2N	70.2W	65	1	40	968	1009	275	MA1	None
5-9/15/1924	1400Z	29.7N	85.3W	75	1	---	980	1011	150	AFL1	AFL1
10-10/21/1924	0100Z	25.8N	81.8W	80	1	20	975	1008	375	BFL1,CFL1	BFL1
4-12/1/1925	0400Z	26.5N	82.2W	65	1	---	(985)	1010	375	BFL1	BFL1
1-7/28/1926	1000Z	29.0N	80.8W	90	2	15	967	1014	300	DFL2,CFL1	DFL2
3-8/25/1926	2300Z	29.2N	90.9W	100	3	20	955	1012	275	LA3	LA3
7-9/18/1926	1200Z	25.7N	80.3W	125	4	20	930	1008	325	CFL4,BFL3	CFL4,BFL3
7-9/20/1926	2200Z	30.3N	87.5W	100	3	15	955	1008	225	AFL3,AL3,MS1	AFL3,AL3
10-10/21/1926*	0300Z	25.0N	80.3W	75	1	20	949	1009	300	BFL1,CFL1	None
None - 1927											
1-8/8/1928	0700Z	27.3N	80.2W	85	2	10	977	1014	150	CFL2	CFL2
4-9/17/1928	0000Z	26.7N	80.0W	125	4	30	929	1009	275	CFL4,BFL3,AFL1,DFL1	CFL4,DFL2
4-9/18/1928	1900Z	32.5N	80.3W	75	1	35	976	1008	350	GA1,SC1	GA1,SC1
1-6/28/1929	2100Z	28.3N	96.4W	80	1	10	982	1007	150	BTX1	BTX1
2-9/28/1929	1300Z	25.0N	80.5W	100	3	30	948	1008	300	BFL3, CFL3	CFL3
2-10/1/1929	0400Z	30.2N	85.7W	70	1	---	975	1011	400	AFL1	AFL2
None - 1930											



Table 4: Major Landfalling (non-continental U.S) Hurricanes - 1921 to 1930. The names listed are unofficial ones that the hurricanes are known by at these locations. Max Winds are the estimated maximum 1-min surface (10 m) winds to occur at along the coast at landfall/closest approach. Saffir-Simpson is the estimated Saffir-Simpson Hurricane Scale at landfall based upon maximum 1-min surface winds. Central Pressure is the minimum central pressure of the hurricane at landfall/closest approach. Central pressure values in parentheses indicate that the value is a simple estimation (based upon a pressure-wind relationship), not directly measured or calculated. Original Winds are the winds in HURDAT that were originally provided at landfall/closest approach.

#/Date	Name	Location	Max Winds	SSHWS	Central Pressure	Original Winds
3-9/11/1921	-----	Dominican Republic	110	3	(951)	80
2-9/16/1922	-----	Barbuda, St. Martin, Anguilla, Anegada	100	3	(961)	90
2-9/21/1922	-----	Bermuda	100	3	970	130
1923 - None						
10-10/19/1924	Huracan sin Precedentes	Cuba	140	5	910	105
1925 - None						
1-7/26/1926	-----	Bahamas	120	4	(938)	120
7-9/17/1926	-----	Bahamas	130	4	(926)	130
10-10/20/1926	-----	Cuba	130	4	934	105
10-10/21/1926	-----	Bahamas	110	3	(948)	100
10-10/22/1926	-----	Bermuda	105	3	962	115
1927 - None						
4-9/12/1928	-----	Guadelope	120	4	940	110
4-9/13/1928	San Felipe	Puerto Rico	140	5	931	140
4-9/15/1928	-----	Bahamas	135	4	(920)	135
2-9/26/1929	-----	Bahamas	125	4	936	125
2-9/3/1930	-----	Dominican Republic	135	4	933	130